

Rate of convergence for a new family of mixed summation-integral type operators

Manoj Kumar

Department of Mathematics and Statistics, Gurukula Kangri Vishwavidyalaya, Haridwar, Uttarakhand, India

Abstract

In the present paper, we introduced a new family of mixed summation integral type operators, to approximate Lebesgue integrable functions on the interval $[0, \infty)$, and study the rate of convergence for these operators. The family of operators is constructed by combing the well known Beta and Baskakov operators. The estimate for rate of convergence is obtained in terms of higher order modulus of continuity, by using the techniques of linear combinations.

Keywords: simultaneous approximation, summation-integral type operators, linear positive operators, modulus of smoothness, linear combination

1. Introduction

Durrmeyer ^[4] introduced the notion of summation-integral type operators to approximate Lebesgue integrable functions on $[0, 1]$. Several researchers proposed and studied rate of convergence for summation-integral type operators ^[1, 2, 3, 6, 8, 9, 12]. Recently, Gupta ^[7] studied rate of convergence of complex Baskako-Szasz-Stancu operators. In the present paper, we introduced a new family of mixed summation integral type operators that is different from the family studied by Gupta ^[7].

Let $C_\gamma[0, \infty)$, $\gamma \geq 0$ be the class of locally integrable functions f defined on $[0, \infty)$ and satisfying the growth condition

$|f(t)| \leq K t^\gamma$, $t \in (0, \infty)$, for some $K > 0$. For $f \in C_\gamma[0, \infty)$ with norm $\|f\|_\gamma = \max \left\{ |f(0)|, \sup_{0 < t < \infty} |f(t)| t^{-\gamma} \right\}$, we

introduce the following family of mixed summation integral type operators as

$$V_n(f, x) = \frac{n-1}{n} \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t) f(t) dt + (1+x)^{-n-1} f(0), \quad x \in [0, \infty), \quad (1.1)$$

$$\text{where } q_{n,v}(x) = \frac{1}{\beta(v+1, n)} x^v (1+x)^{-n-v-1}, \quad p_{n,v}(t) = \binom{n+v-1}{v} \frac{t^v}{(1+t)^{n+v}},$$

and $\beta(v+1, n)$ being the Beta function given by $v!(n-1)/(n+v)!$.

On the other hand, in terms of Dirac delta function $\delta(t)$, the operators (1.1) may be written as

$$V_n(f, x) = \int_0^{\infty} W_n(x, t) f(t) dt, \quad 0 \leq x < \infty, \quad (1.2)$$

where the kernel is given by

$$W_n(x, t) = \frac{n-1}{n} \sum_{v=1}^{\infty} q_{n,v}(x) p_{n,v-1}(t) + (1+x)^{-n-1} \delta(t).$$

Since $f(t) = O(t^\gamma)$ for $f \in C_\gamma[0, \infty)$, and $\int_0^{\infty} t^{r-1} / (1+t)^{s+m} dt$ exists for positive values of r, s only, it follows that for $f \in C_\gamma[0, \infty)$, the operators (1.1) are well defined for each $n > \gamma + 1$.

We may observe that the operators defined by (1.1) are linear positive operators ^[10]. Also the operators V_n reproduce only the constant functions. It is easily verified that the order of approximation by these operators is at best $O(n^{-1})$ howsoever smooth the function may be. For improving the order of approximation, we consider the linear combinations of the operators (1.1) as defined by

$$V_n(f, k, x) = \frac{1}{\Delta} \begin{vmatrix} V_{d_0 n}(f, x) & d_0^{-1} & d_0^{-2} & \dots & d_0^{-k} \\ V_{d_1 n}(f, x) & d_1^{-1} & d_1^{-2} & \dots & d_1^{-k} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ V_{d_k n}(f, x) & d_k^{-1} & d_k^{-2} & \dots & d_k^{-k} \end{vmatrix},$$

where $d_0, d_1, d_2, \dots, d_k$ are $(k + 1)$ arbitrary but fixed distinct positive integers and Δ is the Vandermonde determinant obtained by replacing the operator column of the above determinant by the entries 1. The above linear combinations in alternative form may be rewritten as

$$V_n(f, k, x) = \sum_{j=0}^k C(j, k) V_{d_j n}(f, x) \tag{1.3}$$

where $C(j, k) = \prod_{\substack{i=0 \\ i \neq j}}^k \frac{d_j}{d_j - d_i}$, $k \neq 0$, $C(0, 0) = 1$.

Such type of linear combinations were first considered by May ^[11] to get the better order of approximation for exponential type operators which include Bernstein polynomials, Baskakov operators, Szasz operators, Gauss Weierstrass operators etc. as special cases.

2. Preliminary Results

To prove the main results, we need the following preparatory lemmas and proposition:

Lemma 2.1 For $m \in N^0$ and $x \in [0, \infty)$, if the function $\mu_{n,m}(x)$ be defined by

$$\mu_{n,m}(x) = \frac{n-1}{n} \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t) (t-x)^m dt + (-x)^m (1+x)^{-n-1},$$

then, we have

$$\mu_{n,0}(x) = 1, \quad \mu_{n,1}(x) = \frac{3x}{n-2}, \quad \mu_{n,2}(x) = \frac{x(1+x)(2n-1) + 3x(1+5x)}{(n-2)(n-3)},$$

and there holds the following recurrence relation

$$(n-m-2)\mu_{n,m+1}(x) = x(1+x)[\mu'_{n,m}(x) + 2m\mu_{n,m-1}(x)] + [m(1+2x) + 3x]\mu_{n,m}(x).$$

Furthermore, it follows from the above recurrence relation that

$$\mu_{n,m}(x) = O(n^{-[(m+1)/2]}) \text{ as } n \rightarrow \infty,$$

where $[\alpha]$ denotes the integral part of α .

Proof. The values of $\mu_{n,0}(x), \mu_{n,1}(x)$ easily follow from the definition.

Now, we have

$$\begin{aligned}
 x(1+x)\mu'_{n,v}(x) &= \frac{n-1}{n} \sum_{v=1}^{\infty} x(1+x)q'_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t)(t-x)^m dt \\
 &- m \frac{n-1}{n} \sum_{v=1}^{\infty} x(1+x)q_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t)(t-x)^{m-1} dt \\
 &- [(n+1)(-x)^m(1+x)^{-n-2} + m(-x)^{m-1}(1+x)^{-n-1}]x(1+x)
 \end{aligned}$$

Next using the well known identities

$$x(1+x)q'_{n,v}(x) = [v-(n+1)x]q_{n,v}(x) \quad \text{and} \quad t(1+t)p'_{n,v}(t) = (v-nt)p_{n,v}(t),$$

we obtain

$$\begin{aligned}
 &x(1+x) \left[\mu'_{n,m}(x) + m\mu_{n,m-1}(x) \right] \\
 &= \frac{(n-1)}{n} \sum_{v=1}^{\infty} [v-(n+1)x]q_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t)(t-x)^m dt + (n+1)(-x)^{m+1}(1+x)^{-n-1} \\
 &= \frac{(n-1)}{n} \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} \left[\{(v-1)-nt\} + n(t-x) + (1-x) \right] p_{n,v-1}(t)(t-x)^m dt + (n+1)(-x)^{m+1}(1+x)^{-n-1} \\
 &= \frac{(n-1)}{n} \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} t(1+t)p'_{n,v-1}(t)(t-x)^m dt + \frac{(n-1)}{n} n \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t)(t-x)^{m+1} dt \\
 &\quad + \frac{(n-1)}{n} (1-x) \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t)(t-x)^m dt + (n+1)(-x)^{m+1}(1+x)^{-n-1} \\
 &= \frac{(n-1)}{n} \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} \left[(1+2x)(t-x) + (t-x)^2 + x(1+x) \right] p'_{n,v-1}(t)(t-x)^m dt \\
 &\quad + n\mu_{n,m+1}(x) + \frac{(n-1)}{n} (1-x) \sum_{v=1}^{\infty} q_{n,v}(x) \int_0^{\infty} p_{n,v-1}(t)(t-x)^m dt + (-x)^{m+1}(1+x)^{-n-1} \\
 &= -[m(1+2x) + 3x]\mu_{n,m}(x) + (n-m-2)\mu_{n,m+1}(x) - mx(1+x)\mu_{n,m-1}(x)
 \end{aligned}$$

This completes the proof of recurrence relation.

Corollary 2.2^[5]. Let γ and δ be two positive numbers, then for any $m > 0$ there exists a constant C such that

$$\left\| \int_{|t-x| \geq \delta} W_n(x,t)t^\gamma dt \right\|_{C[a,b]} \leq C n^{-m}.$$

Proposition 2.3^[5]. For $m \in N$ and sufficiently large $n \in N$, there holds the following relation

$$V_n((t-x)^m, k, x) = n^{-(k+1)} [Q(m, k, x) + o(1)],$$

where $Q(m, k, x)$ are certain polynomials in x of degree at most m .

Lemma 2.4 ⁽¹⁵⁾. For $0 < a < a_1 < a_2 < b_2 < b_1 < b < \infty$ and sufficiently small $\delta > 0$, let the $(2k+2)$ th order Steklov mean $g_{2k+2,\delta}(t)$ corresponding to $g \in C_\gamma[0, \infty)$ be defined by

$$g_{2k+2,\delta}(t) = \delta^{-(2k+2)m} \int_{-\delta/2}^{\delta/2} \int_{-\delta/2}^{\delta/2} \dots \int_{-\delta/2}^{\delta/2} [g(t) - \Delta_\eta^{2k+2} g(t)] \prod_{i=1}^{2k+2} dt_i$$

where $\eta = \frac{1}{2k+2} \sum_{i=1}^{2k+2} t_i$, $t \in [a, b]$ and $\Delta_\eta^{2k+2} g(t)$ is the $(2k+2)$ th forward difference with step length η . Then we have

- i) $g_{2k+2,\delta}$ has continuous derivatives up to order $(2k+2)$ on $[a, b]$;
- ii) $\|g_{2k+2,\delta}^{(r)}\|_{C[a_1, b_1]} \leq C_1 \delta^{-r} \omega_r(g, \delta, a_1, b_1)$, $r = 1, 2, 3, \dots, (2k+2)$;
- iii) $\|g - g_{2k+2,\delta}\|_{C[a_2, b_2]} \leq C_2 \omega_{2k+2}(g, \delta, a_1, b_1)$;
- iv) $\|g_{2k+2,\delta}\|_{C[a_2, b_2]} \leq C_3 \|g\|_\gamma$,

where C_1, C_2, C_3 are certain unrelated constants independent of g and δ .

Main Results

In this section we state and prove the following direct results

Theorem 3.1. If $(2k+2)$ th derivatives of $f \in C_\gamma[0, \infty)$ exist at a point $x \in (0, \infty)$, then we have

$$n^{k+1} [V_n(f, k, x) - f(x)] = \sum_{i=k+1}^{2k+2} \frac{f^{(i)}(x)}{i!} Q(i, k, x) + o(1) \tag{3.1}$$

and

$$\lim_{n \rightarrow \infty} n^{k+1} [V_n(f, k+1, x) - f(x)] = 0 \tag{3.2}$$

where $Q(i, k, x)$ are certain polynomials in x of degree i .

Proof. By Taylor’s finite expansion of $f(t)$ we have

$$f(t) = \sum_{i=0}^{2k+2} \frac{f^{(i)}(x)}{i!} (t-x)^i + \varepsilon(t, x) (t-x)^{2k+2}, \text{ where } \varepsilon(t, x) \rightarrow 0 \text{ as } t \rightarrow x.$$

Now,

$$\begin{aligned} & n^{k+1} [V_n(f, k, x) - f(x)] \\ &= n^{k+1} \sum_{i=1}^{2k+1} \frac{f^{(i)}(x)}{i!} V_n((t-x)^i, k, x) + n^{k+1} \sum_{j=0}^k C(j, k) V_{d_j n}(\varepsilon(t, x) (t-x)^{2k+2}, x) \\ &= E_1 + E_2 \text{ (say)} \end{aligned}$$

Since

$$\mu_{d_j n, i}(x) = \frac{P_0(x)}{(d_j n)^{\lceil (i+1)/2 \rceil}} + \frac{P_1(x)}{(d_j n)^{\lceil (i+1)/2 \rceil + 1}} + \dots + \frac{P_{\lfloor i/2 \rfloor}(x)}{(d_j n)^i},$$

where $P_i(x), i = 0, 1, 2, \dots, \lfloor i/2 \rfloor$ are certain polynomials in x of degree at most $\lfloor i/2 \rfloor$. Therefore, by Proposition 2.3, we get

$$\sum_{j=0}^k C(j, k) \mu_{d_j n, i}(x) = \frac{1}{\Delta} \begin{vmatrix} \frac{P_0(x)}{(d_0 n)^{\lceil (i+1)/2 \rceil}} + \frac{P_1(x)}{(d_0 n)^{\lceil (i+1)/2 \rceil + 1}} + \dots + \frac{P_{\lfloor i/2 \rfloor}(x)}{(d_0 n)^i} & d_0^{-1} & d_0^{-2} & \dots & d_0^{-k} \\ \frac{P_0(x)}{(d_1 n)^{\lceil (i+1)/2 \rceil}} + \frac{P_1(x)}{(d_1 n)^{\lceil (i+1)/2 \rceil + 1}} + \dots + \frac{P_{\lfloor i/2 \rfloor}(x)}{(d_1 n)^i} & d_1^{-1} & d_1^{-2} & \dots & d_1^{-k} \\ \dots & \dots & \dots & \dots & \dots \\ \frac{P_0(x)}{(d_k n)^{\lceil (i+1)/2 \rceil}} + \frac{P_1(x)}{(d_k n)^{\lceil (i+1)/2 \rceil + 1}} + \dots + \frac{P_{\lfloor i/2 \rfloor}(x)}{(d_k n)^i} & d_k^{-1} & d_k^{-2} & \dots & d_k^{-k} \end{vmatrix}$$

$$= n^{-(k+1)} [Q(i, k, x) + o(1)], \quad i = (k+1), (k+2), \dots, (2k+2).$$

where Δ is the Vandermonde determinant obtained by replacing the first column by the entries 1. Consequently, we get

$$E_1 = n^{k+1} \sum_{i=1}^{2k+2} \frac{f(x)}{i!} \sum_{j=0}^k C(j, k) \mu_{d_j n, i}(x) = \sum_{i=k+1}^{2k+2} \frac{f^{(i)}(x)}{i!} Q(i, k, x) + o(1).$$

To prove the assertion (3.1), it is sufficient to show that $E_2 \rightarrow 0$ as $n \rightarrow \infty$.

For a given $\varepsilon > 0$ there exists a $\delta > 0$ such that for $0 < |t - x| < \delta, |\varepsilon(t, x)| < \varepsilon$ and for

$$|t - x| \geq \delta, |\varepsilon(t, x)|(t - x)^{2k+2} \leq K t^\gamma, \text{ for some positive constant } K.$$

If $\phi_\delta(t)$ is the characteristic function of the interval $(x - \delta, x + \delta)$, then we have

$$|E_2| \leq n^{k+1} \sum_{j=0}^k C(j, k) \left[V_{d_j n}(|\varepsilon(t, x)|(t - x)^{2k+2} \phi_\delta(t), x) + V_{d_j n}(|\varepsilon(t, x)|(t - x)^{2k+2} [1 - \phi_\delta(t)], x) \right]$$

$$= E_{21} + E_{22} \text{ (say)}$$

Applying Lemma 2.1 and Corollary 2.2, we obtain

$$|E_{21}| \leq \varepsilon n^{k+1} \sum_{j=0}^k |C(j, k)| \max_{0 \leq j \leq k} \mu_{d_j n, 2k+2}(x) \leq C_1 \varepsilon$$

and

$$|E_{22}| \leq \varepsilon n^{k+1} \sum_{j=0}^k |C(j, k)| \int_{|t-x| \geq \delta} W_{d_j n}(x, t) |\varepsilon(t, x)| (t - x)^{2k+2} dt$$

$$\leq n^{k+1} \sum_{j=0}^k |C(j, k)| \int_{|t-x| \geq \delta} W_{d_j n}(x, t) t^\gamma dt \leq C_2 n^{k+1} \sum_{j=0}^k |C(j, k)| n^{-m} = O(1)$$

for some $m \geq (k + 1)$.

Finally, combining the estimates of E_{21} and E_{22} , we get (3.1). The proof of (3.2) follows easily along the lines of the proof of (3.1), by noticing the fact that

$$V_n((t - x)^i, k + 1, x) = O(n^{-(k+2)}).$$

This completes the proof.

Theorem 3.2. Let $f \in C_r[0, \infty)$. If $f^{(p)}, 1 \leq p \leq 2k + 2$ exists and is continuous with the modulus of continuity $\omega(f^{(p)}, \delta)$ on $[0, a]$. Then for sufficiently large $n \in N$, we have

$$\|V_n(f, k, \cdot) - f\|_{C[0,b]} \leq \max \{C_1 n^{-p/2} \omega(f^{(p)}, n^{-1/2}), C_2 n^{-(k+1)}\}$$

where $C_1 = C_1(k, p), C_2 = C_2(k, p, f)$ and $0 < b < a$.

Proof. For all $t > 0$ and $x \in [0, a]$, we have

$$f(t) = \sum_{i=0}^p \frac{f^{(i)}(x)}{i!} (t - x)^i + [f^{(p)}(\xi) - f^{(p)}(x)] \frac{(t - x)^p}{p!} + h(t, x) \chi(t), \tag{3.3}$$

where ξ lies between t and x and $\chi(t)$ is the characteristic function of the set $(0, \infty) \setminus [0, a]$. The function $h(t, x)$ is bounded by $K t^\gamma |t - x|^p, K > 0$.

Using (3.3), we have

$$\begin{aligned} V_n(f, k, x) &= \sum_{j=0}^k C(j, k) V_{d,n}(f, x) = \sum_{j=0}^k C(j, k) \int_0^\infty W_{d,n}(x, t) f(t) dt \\ &= \sum_{j=0}^k C(j, k) \int_0^\infty W_{d,n}(x, t) \left[\sum_{i=0}^p \frac{f^{(i)}(x)}{i!} (t - x)^i + [f^{(p)}(\xi) - f^{(p)}(x)] \frac{(t - x)^p}{p!} + h(t, x) \chi(t) \right] dt \\ &= \sum_{i=0}^p \frac{f^{(i)}(x)}{i!} \sum_{j=0}^k C(j, k) \int_0^\infty W_{d,n}(x, t) (t - x)^i dt \\ &\quad + \sum_{j=0}^k C(j, k) \int_0^\infty W_{d,n}(x, t) [f^{(p)}(\xi) - f^{(p)}(x)] \frac{(t - x)^p}{p!} dt + \sum_{j=0}^k C(j, k) \int_0^\infty W_{d,n}(x, t) h(t, x) \chi(t) dt \\ &= J_1 + J_2 + J_3 \text{ (say)} \end{aligned}$$

Making use of Lemma 2.1, we get

$$J_1 = f(x) + o(n^{-(k+1)}) \text{ uniformly for all } x \in [0, a]$$

Since for all $\delta > 0$, we have

$$|f^{(p)}(\xi) - f^{(p)}(x)| \leq \omega(f^{(p)}, |\xi - x|) \leq \omega(f^{(p)}, |t - x|) \leq \left[1 + \frac{|t - x|}{\delta} \right] \omega(f^{(p)}, \delta).$$

Therefore, we have

$$|J_2| \leq \frac{\omega(f^{(p)}, \delta)}{p!} \sum_{j=0}^k |C(j, k)| \int_0^\infty W_{d_j n}(x, t) \left[|t-x|^p + \frac{|t-x|^{p+1}}{\delta} \right] dt$$

Using Schwarz inequality and Lemma 3.1, we get

$$|J_2| \leq \frac{\omega(f^{(p)}, \delta)}{p!} \sum_{j=0}^k |C(j, k)| \left[O((d_j n)^{-p}) + \frac{1}{\delta} O((d_j n)^{-(p+1)}) \right]^{1/2}$$

Consequently, choosing $\delta = n^{-1/2}$, we obtain

$$|J_2| \leq C_3 n^{-p/2} \omega(f^{(p)}, n^{-1/2})$$

Next, choosing a positive number s such that $|t-x| \geq s$, we get

$$|J_3| \leq \sum_{j=0}^k |C(j, k)| \int_{|t-x| \geq s} W_{d_j n}(x, t) t^\gamma |t-x|^p dt.$$

Applying Schwarz inequality and Corollary 2.2, we have

$$\begin{aligned} |J_3| &\leq \sum_{j=0}^k |C(j, k)| \left[\int_{|t-x| \geq s} W_{d_j n}(x, t) e^{2\gamma t} dt \right]^{1/2} \left[\int_{|t-x| \geq s} W_{d_j n}(x, t) (t-x)^{2p} dt \right]^{1/2} \\ &\leq \sum_{j=0}^k |C(j, k)| (Kn^{-m})^{-1/2} O((d_j n)^{-p/2}) \leq C_4 n^{-(k+1)}. \end{aligned}$$

Finally, combining the estimates of J_1, J_2 and J_3 , the Theorem 3.2 follows.

This completes the proof.

Our next result is the rate of convergence in terms of higher order modulus of continuity, which is stated as

Theorem 3.3. Let $f \in C_\gamma[0, \infty)$ and $0 < a_1 < a_2 < b_2 < b_1 < \infty$. Then for sufficiently large $n \in \mathbb{N}$, we have

$$\|V_n(f, k, \cdot) - f\|_{C[a_2, b_2]} \leq K \left[n^{-(k+1)} \|f\|_\gamma + \omega_{2k+2}(f, n^{-1/2}; a_1, b_1) \right],$$

where the constant K is independent of f and n .

Proof. First by linearity property of the operators (1.3), we have

$$\begin{aligned} \|V_n(f, k, \cdot) - f\|_{C[a_2, b_2]} &\leq \|V_n((f - f_{2k+2, \delta}), k, \cdot)\|_{C[a_2, b_2]} \\ &\quad + \|V_n(f_{2k+2, \delta}, k, \cdot) - f_{2k+2, \delta}\|_{C[a_2, b_2]} + \|f - f_{2k+2, \delta}\|_{C[a_2, b_2]} \\ &= A_1 + A_2 + A_3 \text{ (say)} \end{aligned}$$

By property (iii) of Lemma 2.4, we have

$$A_3 \leq C_1 \omega_{2k+2}(f^{(r)}, \delta; a_1, b_1)$$

Next by Theorem 3.1, we have

$$A_2 \leq C_2 n^{-(k+1)} \sum_{j=k+1}^{2k+2} \|f_{2k+2,\delta}^{(j)}\|_{C[a_1,b_1]}$$

By using the interpolation property due to Goldberg and Meir^[5] for each $j = k + 1, k + 2, \dots, 2k + 2$, we have

$$\|f_{2k+2,\delta}^{(j)}\|_{C[a_1,b_1]} \leq C_3 \left[\|f_{2k+2,\delta}\|_{C[a_1,b_1]} + \|f_{2k+2,\delta}^{(2k+2)}\|_{C[a_1,b_1]} \right] \leq C_4 \left[\|f_{2k+2,\delta}^{(2k+2)}\|_{C[a_1,b_1]} + \|f\|_\gamma \right]$$

Again using Lemma 2.4, it follows that

$$\begin{aligned} A_1 &= \left\| V_n \left((f - f_{2k+2,\delta}), k, \cdot \right) \right\|_{C[a_2,b_2]} \\ &\leq \left\| \sum_{j=0}^k |C(j,k)| \int_0^\infty W_{d,n}(x,t) |f(t) - f(x)| dt \right\|_{C[a_2,b_2]} \\ &\leq \left\| \sum_{j=0}^k |C(j,k)| \left[\int_{|t-x| \leq \delta} + \int_{|t-x| > \delta} \right] W_{d,n}(t,x) |f(t) - f(x)| dt \right\|_{C[a_2,b_2]} \\ &\leq \|f - f_{2k+2,\delta}\|_{C[a_2-\delta,b_2+\delta]} + C_5 n^{-m} \|f\|_\gamma \\ &\leq C_6 \omega_{2k+2}(f, \delta; a_1, b_1) + C_5 n^{-m} \|f\|_\gamma \end{aligned}$$

Finally, choosing $\delta = n^{-1/2}$, and combining the estimates of A_1, A_2 and A_3 , we get the desired result. This completes the proof.

References

1. Acar T, Gupta V, Aral A. Rate of convergence for generalized Szasz operators, Bulletin of Math Sci., 2011; 1:99-113.
2. Aniol G, Taberska PP. On the rate of convergence of the Durrmeyer polynomials, Ann. Soc. Math. Pol., Ser, I, Commentant Math., 1990; 30:9-17.
3. Cheng F. On the rate of convergence of Bernstein polynomials of functions of bounded variation, J Approx. Theory. 1983; 39:259-274.
4. Durrmeyer JL, Une Formule d' Inversion de la Transformee de Laplace, Applications a la Theory des Moments, These de 3e cycle, Faculte des Sciences del' Universite' de Paris, 1967.
5. Goldberg S, Meir V. Minimum moduli of ordinary differential operators, Proc. London Math. Soc. 1971; 23:1-15.
6. Govil NK, Gupta V. Rate of convergence for Baskakov type operators on functions of bounded variation, Journal of Combinatorics, Information and System Sciences. 2006; 31(1-4):1-17.
7. Gupta V. Over convergence of Complex Baskako-Szasz-Stancu Operators, Med J. Math., 2015; 12:455-470.
8. Gupta V, Yadav R. Rate of convergence for generalized Baskakov operators, Arab J. Mathematical Sciences. 2012; 18(1):39-50.
9. Karsi H, Gupta V. Rate of Convergence by Nonlinear Integral Operators for Functions of Bounded Variation, Calcolo. 2008; 45:87-97.
10. Korovkin PP. Linear Operators and Approximation Theory, Hindustan Publ. Corp. Delhi, the Russian original appeared in 1959-1960.
11. May CP. Saturation and inverse theorems for combinations of a class of exponential type operators, Canad. J Math., 1976; 28:1224-1250.
12. Zeng XM, Gupta V. Rate of convergence of Baskakov-Bézier type operators for locally bounded functions, Comput. Math. Appl., 2002; 44(10-11):1445-1453.