

Phytochemical analysis of leaves and tubers of *Gloriosa superba* L. Endangered plant

Aradhana Mishra¹, Neeta Singh²

¹ Research Scholar, Department of Botany, Govt. Science P.G. College, (A.P.S. University), Rewa, Madhya Pradesh, India

² Professor, Department of Botany, Govt. Girls P.G. College, Rewa, Madhya Pradesh, India

Abstract

Gloriosa superba L. (Family: Liliaceae) is one of the endangered species among the medicinal plants which is a striking tuberous climbing plant with brilliant wavy-edged yellow and red flowers common in forest throughout India. The objective of this study was to evaluate the phytoconstituents of *Gloriosa superba* leaf and tuber extracts. All experiments were conducted following standard procedures. The results of the various phytochemical tests indicated that the plant to be rich in various biologically active compounds which could serve as potential source of the crude drugs that can be used as a complementary source of traditional medicines.

Keywords: *Gloriosa superba*, endangered, phytoconstituents, leaves and tubers

1. Introduction

Gloriosa superba Linn. Of family Liliaceae, is one of the endangered species among the medicinal plants (Badola, 2002) [1] which is a striking tuberous climbing plant with brilliant wavy-edged yellow and red flowers that appears from November to March every year (Rajak and Rai, 1990) [2]. Herbaceous tendril climber; root stock tuberous, naked, stem sparingly branched; leaves sessile or nearly so, flowers axillary, solitary, at first greenish, becoming yellow and finally red.

The tuberous root stocks of glory lily, boiled with Sesamum oil is applied twice a day on the joints, affected with arthritis reduces pain (Singh, 1993) [3]. Seeds are used for relieving rheumatic pain and as a muscle relaxant (Nadkarni, 2002) [4]. The tuber is useful in itching, skin diseases including wounds and ailments caused by vitiated kapha and vata, can be administered to a delivered mother along with spirituous drink to give relieve to her postnatal complaints. It is pungent, thermogenic, and used as a purgative. In the world market, the tubers are considered as rich sources of colchicines and gloriosine (Srivatava and Chandra, 1977 and Prajapati *et al.* 1977) [5, 6]. Roots are acrid, anthelmintic, antipyretic, bitter, digestive, expectorant, highly poisonous and promoting expulsion of the placenta. Root paste is effective against paralysis, rheumatism, snake bite and insect bites.

It is one of the major medicinal plants in India cultivated for its seeds which are exported to developed countries for pharmaceutical use. However, not much is known about the chemical composition of the plant leaves and tubers (Chitra and Rajamani, 2009) [7].

The aim of this study was to determine the phytochemical properties of *Gloriosa superba* leaves and tubers with the aim of further isolation and characterization and the possible use of these substances in industries and medicinal science.

2. Materials and Methods

Plant materials: *Gloriosa superba* leaves and tubers were collected from forest of Govindgarh district Rewa (M.P.) India and Identified by Dr. Sanjeev Dubey, Prof. & Head,

Govt. Science P.G. College, Rewa and has been deposited the herbarium sheets in Bot. Deptt. Research center.

Preparation of plant extract: The acetone, chloroform, dichloromethane and methanol extracts of the leaves and tubers of the plant were prepared according to standard methods (Harborne, 1973) [8]. The plant parts such as leaves and tubers were collected and shade dried for about two weeks and ground into coarse powder. About 50 g powder of each plant part was separately extracted with 125 ml of acetone using soxhlet apparatus. The same powders were also extracted with chloroform, dichloromethane and methanol. The extracts were concentrated to dryness to yield crude residue. These residues were used for preliminary phytochemical screening of secondary metabolites.

Table 1: Preliminary phytochemical tests for the presence of active constituents in *Gloriosa superba* leaves

S. No.	Compounds	Solvents used			
		Acetone	Chloroform	Dichloro methane	Methanol
1.	Alkaloids	-	-	-	-
2.	Flavonoids	+	+	+	+++
3.	Saponins	-	-	-	-
4.	Phenols	+++	++	++	+++
5.	Steroids	+	+	+	++
6.	Glycosides	+++	+++	+++	+++
7.	Tannins	+++	+	+	+++
8.	Resins	-	-	-	-

Phytochemical screening

The leaves and tubers extract of *Gloriosa superba* were analyzed for the presence of alkaloids, flavonoids, saponins, glycosides, steroids, phenols, tannins and resins according to standard methods (Harborne, 1973) [8].

Screening for Alkaloids

Meyer's reagent (potassium mercuric iodide) 1.36 gm of mercuric chloride was dissolved in 60 ml of distilled water and 5 gm of potassium iodide was dissolved in 10 ml of water. These two solutions were mixed and diluted to 100

ml with distilled water. To 1 ml of the extract, a few drops of reagent were added. Formation of white or pale precipitate showed the presence of alkaloids.

Screening for Flavonoids

In a test tube containing 0.5 ml of extract, 5 to 10 drops of diluted HCl and small piece of ZnCl or magnesium were added and the solution was boiled for few minutes. The appearance of reddish pink or dirty brown colour indicates the presence of flavonoids.

Screening for Saponins

In a test tube containing about 5 ml of the extract, few drops of sodium bicarbonate was added. The mixture was shaken vigorously and kept for 3 minutes. A honeycomb like froth was formed and it showed the presence of saponins.

Screening for Glycosides

A small amount of extract was dissolved in 1 ml of water and aqueous sodium hydroxide solution was added. Formation of yellow colour indicates the presence of glycosides.

Screening for Steroids

To 2.0 ml of extract, 1.0 ml of concentrated sulphuric acid was added carefully along the sides of the test tube. A red colour produced in the chloroform layer shows the presence of steroids.

Screening for Phenols

Ferric chloride test: To 1 ml of the extract 3 ml of distilled water followed by few drops of 10% aqueous Ferric chloride solution was added. Formation of blue or green colour indicates the presence of phenols

Screening for Tannins

Lead acetate test: In a test tube containing about 5 ml of the extract, a few drops of 1% solution of lead acetate was added. A yellow or red precipitate indicates the presence of tannins.

Screening for Resins

To 2.0 ml of extract 5 ml of acetic anhydride was added, dissolved by gently heating, cooling and then 0.5 ml of sulphuric acid was added. Bright purple colour indicates the presence of resins.

3. Results

The results of preliminary phytochemical screening of acetone, chloroform, dichloromethane and methanol leaf and tuber extracts of *Gloriosa superba* are presented in Table 1 and 2. All the leaves and tuber extracts showed abundant occurrence of phenols, tannins and glycosides. Alkaloids were found to be present in higher concentration in all the extracts of the tubers but absent in the leaves extracts. Flavonoids and steroids were found in a moderate concentration in all the extracts of leaves and tubers. Saponins were found in moderate concentration in chloroform, dichloromethane, methanol tuber extracts but showed the absence in the acetone extract of tubers and all the extracts of leaves. Resins were absent in all the extracts of the two plant portions namely the leaves and tubers.

4. Discussion

The interest in the study of medicinal plants as a source of pharmacologically active compounds has increased worldwide (Rios and Recio, 2005) [9]. In most developing countries of the world, plants are the main medicinal sources used in treating infectious diseases (Sofowora, 1993) [10].

The various phytochemical compounds detected are known to exhibit medicinal activity as well as physiological activity (Sofowora, 1993) [10].

There are records that show the benefits of these compounds detected from *Gloriosa superba*. For example: Many of the earliest isolated pure compounds with biological activity were alkaloids. Naturally occurring alkaloids are nitrogenous compounds that constitute the basic active principles of flowering plants. Alkaloids are formed as metabolic by-products and have been reported to be responsible for the antibacterial activity (Doughari, 2006) [11]. Phenolics are the largest group of and have been said to account for most of the antioxidant activity of plant extracts (Thabrew *et al.* 1998) [12]. The trace quantities of phenolic compounds help to prevent the death of the crop, since phenolic compounds from plant extracts act as antimicrobial agent (Ofokansi *et al.* 2005) [13]. Phenolics and alkaloids detected in the extracts are compounds that have been documented to possess medicinal properties and health-promoting effects (Salah *et al.* 1995, Okwu. 2004, Liu, 2004) [14-16]. Glycosides are nonvolatile and lack fragrance. Glycosides serve as defense mechanisms against predation by many microorganisms, insects and herbivores (De M *et al.* 1999) [17].

Table 2: Preliminary phytochemical tests for the presence of active constituents in *Gloriosa superba* tubers

S. No	Compounds	Solvents used			
		Acetone	Chloroform	Dichloro Methane	Methanol
1.	Alkaloids	+++	+++	+++	+++
2.	Flavonoids	++	+	+	++
3.	Saponins	-	++	+	+++
4.	Phenols	+++	+	++	+++
5.	Steroids	++	++	+	++
6.	Glycosides	+++	+++	+++	+++
7.	Tannins	++	++	++	++
8.	Resins	-	-	-	-

Steroidal compounds are of importance and interest in pharmacy due to their relationship with such compounds as sex hormones. Plant steroids are known to be important for their cardiostimulant activities, they possess insecticidal and anti-microbial properties. They are routinely used in medicine because of their profound biological activities (Okwu, 2001 and Denwick *et al.* 2002) [18-19].

Saponins exhibit cytotoxic effect and growth inhibition against a variety of cell making them have anti-inflammatory and anticancer properties. They also show tumor inhibiting activity on animals (Akindalunsi and Salawu, 2005) [20]. Plant saponins help humans to fight fungal infections, combat microbes and viruses, boost the effectiveness of certain vaccines and knock out some kinds of tumor cells, particularly lung and blood cancers (Barkat *et al.* 1995) [21]. These compounds served as natural antibiotics, which help the body to fight infections and

microbial invasion (Sodipo *et al.* 2000) ^[22]. Tannins have been traditionally used for protection of inflamed surfaces of the mouth and treatment of catarrh, wounds, hemorrhoids and diarrhea (Ogunleye and Ibitoye, 2003) ^[23]. Plant tannins have been recognized for their pharmacological properties and are known to make trees and shrubs (Heslem, 1989) ^[24]. Flavonoids are widely distributed group of polyphenolic compounds, characterized by a common benzopyrene ring structure. The biological functions of flavonoids apart from its antioxidant properties include protection against allergies, inflammation, free radicals, platelet aggregation, microbes, ulcers, hepatoxins, viruses and tumors (Okwu and Omodamiro, 2005) ^[25]. Flavonoids reduced cancers by interfering with the enzymes that produce estrogen (Okwu and Omodamiro, 2005 and Farquer, 1990) ^[25-26].

5. Conclusion

In view of all these various uses associated with these compounds found in *Gloriosa superba* leaves and tubers extracts, we recommend further research on this plant to quantify the concentration of these compounds per known amount for industrial use. We believe these compounds in *Gloriosa superba* leaves and tubers could be harnessed for industrial and medicinal sciences utilization.

6. Acknowledgement

The authors are greatly indebted to Principal and Head of Botany Deptt. Of Govt. Science College, Rewa (M.P.) who permitted to carry out this work.

7. References

1. Badola HK. Endangered medicinal plant species in Himachal Pradesh. A report on the International Workshop on "Endangered Medicinal Plant Species in Himachal Pradesh", organized by G.B. Pant Institute of Himalayan Environment and Development at Himachal Unit, Mohal-Kullu during 18-19 March 2002. *Curr. Sci.* 2002; 83:797-798.
2. Rajak RC, MK Rai, *Herbal Medicines Biodiversity and Conservation Strategies*. International Book Distributors, 1990, 75-79.
3. Singh VK. Selected Indian Folk medicinal claims and their relevance in primary health care programme. *Glimpses Plant Res.* 1993; 10:147-152.
4. Nadkarni AK. *Indian Materia Medica*. Popular Prakashan limited, 2002, 579-581.
5. Srivastava UC, Chandra V. *Gloriosa superba* Linn. (Kalihari) - An important colchicine producing plant. *J. Res. Indian Med.* 1977; 10:92-95.
6. Prajapati ND, Purohit SS, Sharma AK, Kumar T. *A hand book of Medicinal Plants*: Agro bios publishers (India); Jodhpur, 2003, 221-234.
7. Chitra R, Rajamani K. Per Se Performance and Correlation for Yield and its Quality Characters in Glory Lily (*Gloriosa superba*, L.). *Academic Journal of Plant Sciences.* 2009; 2(1):39-43.
8. Harborne JB. *Phytochemical methods*. Chapman and Hall Ltd., London, 1973, 49-188.
9. Rios JL, Recio MC. Medicinal plants and antimicrobial activity. *J. Ethno pharmacology.* 2005; 100:80-84.
10. Sofowora LA. *Medicinal plants and traditional medicine in Africa*. Spectrum Books Ltd, Ibadan, 1993, 55-71.
11. Doughari JH. Antimicrobial activity of *Tamarindus indica* Linn. *Trop J. Pharm. Res.* 2006; 5:597.
12. Thabrew MI, Hughes RD, McFarlane IG. Antioxidant activity of *Osbeckia aspera*. *Phytother. Res.* 1998; 12:288-290.
13. Ofokansi KC, Esimone CO, Anele CK. Evaluation of the *in vitro* combined anti-bacterial effects of the leaf extracts of *Bryophyllum pinnatum* (FAM Crassulaceae) and *Ocimum gratissimum* (Fam Labiatae). *Plant Prod. Res. J.* 2005; 9:23-27.
14. Salah W, Miller NJ, Pagauga G, Tijburg, Bolwell GP, Rice E, Evans C. Polyphenolic flavonols as scavenger of aqueous phase radicals and chain breaking antioxidants. *Arch. Biochem. Biol.* 1995; 2: 339-346.
15. Okwu DE. Phytochemical and vitamin content of indigenous spices of South Eastern Nigeria. *J. Sustain. Agric. Environ.* 2004; 6:30-34.
16. Liu RH. Potential synergy of phytochemicals in cancer prevention: mechanism of action. *J. Nutr.* 2004; 134:3479S-3485S.
17. De M, Krishina De A, Banerjee AB. Antimicrobial screening of some Indian spices. *Phytother. Res.* 1999; 13:616-618.
18. Okwu DE. Evaluation of the chemical composition of indigenous spices and flavouring agents. *Global J. Pure Appl. Sci.* 2001; 7:455-459.
19. Denwick PM. *Natural products: A Biosynthetic Approach*. 2nd Edn. England. John Wiley and Sons, Ltd, 2002, 241-243.
20. Akindahunsi AA, Salawu SO. Phytochemical screening and nutrient-antinutrient composition of selected tropical green leafy vegetables. *Afr. J. Biotech.* 2005; 4:497-501.
21. Barakat MZ, Shahab SK, Darwin N, Zahemy EI. Determination of ascorbic acid from plants. *Anal. Biochem.* 1995; 53:225-245.
22. Sodipo OA, Akiniyi JA, Ogunbanosu. Studies on certain characteristics of extracts of bark of *Pansinystalia macrucas* (K. Schem) Piere. Exbeile. *Global J. Pure Appl. Sci.* 2000; 6:83-87.
23. Ogunleye DS, Ibitoye SF. Studies of antimicrobial activity and chemical constituents of *Ximenia Americana*. *Trop. J. Pharm. Res.* 2003; 2(2):239-241.
24. Heslem E. *Plant Polyphenol: Vegetal tannin related-Chemistry and Pharmacology and Natural Products* Cambridge University Press, 1989, 169.
25. Okwu DE, Omodamiro OD. Effects of hexane extract and phytochemical content of *Xylopiya aethiopia* and *Ocimum gratissimum* on the uterus of guinea pig. *Biol. Res.* 2005, 3.
26. Farquer JN. *Plant Sterols. Their biological effects in humans*. Handbook of lipids in human nutrition. BOCA Raton FL CRC Press, 1990, 101-105.